

Léo Valette*, Julian Pietragalla, Marie-Angélique Laporte, Agbona Afolabi, Ousmane Boukar, Steven B. Cannon, Brian W. Diers, Kate A. Dreher, Pooran M. Gaur, Alberto Fabio Guerrero, C. Tom Hash, Vilma Hualla, Drabo Inoussa, Scott R. Kalberer, Clarisse Kondombo-Barro, Shiv Kumar, Antonio Lopez-Montes, Naama Menda, Randall Nelson, Sam Ofodile, Suyash Patil, Peteti Prasad, Karthika Rajendran, Jean-Francois Rami, Abhishek Rathore, Ruairadh Sackville Hamilton, Rosemary Shrestha, Reinhard Simon, Niaba Teme, Eva Weltzien-Rattunde and Elizabeth Arnaud*.

* Bioversity International, Montpellier, France. Email: e.arnaud@cgiar.org

The Crop Ontology development

Trait Dictionary template

- Developed by the Crop Ontology and the Integrated Breeding Platform to **support the creation of ontologies**
- Populated by breeders with their **traits**, observation **methods** and reporting **scales**



The **Breeding Management System** uses the Trait Dictionaries to:

- Create breeders' fieldbook
- Annotate and store breeders' data <http://integratedbreeding.net/>



The **Crop Ontology online tool** features:

- The **publication** of ontologies from the Trait Dictionary template or OBO files
- The **browsing** of term definitions and relations
- An **Application Programming Interface** to provide databases and web applications with ontologies in Excel, OBO, RDF, JSON formats

<http://www.cropontology.org>

(<https://github.com/bioversity/Crop-Ontology>)

Improvement in phenotype annotation

To annotate phenotypes, Crop Ontology supplies breeders with:

- Traits** i.e. the observed plant entities (e.g. leaf, grain) and attributes (e.g. color, weight)
- Methods** i.e. the protocols to observe the trait
- Scales** i.e. the units or categories that can express the trait observation


Thus, a plant phenotype had to be annotated with 3 identifiers for the trait, the method and the scale, respectively. Yet, breeders' fieldbooks and phenotype databases are often designed to **annotate a datapoint with only one identifier**. In May 2015, Crop Ontology has consequently been revised to integrate the variables.

1 **Variable** = {1 Trait, 1 Method, 1 Scale}

Examples of variables:

Variable	Trait	Method	Scale
PltHt_Meas_cm	Plant height i.e., the distance from ground level to the tip of the spike	Measure the height of a plant with a ruler	cm
PltHt_Av_cm		Compute the average of 5 to 10 individual plant height measurements	cm
PltHt_Est_0to5		Visually estimate the average plant height	Score
GW100_Meas_g	100 grain weight i.e., the weight of 100 dehulled grains	Weigh a defined number of grains. Then, divide the weight by the number of grains and multiply by 100	g per 100 grains

CO schema upgrade



Online visualization

Language of submission EN

Variable ID CO_339:0000254


Variable name GrPrtCent_Comp_Ptc

Context of use Evaluation trials

Variable status Recommended

<http://www.cropontology.org>

Crop Trait Dictionary upgrade



Next Steps

- Complete the upgrade of the Trait Dictionaries
- Start upgrading the Trait Dictionaries of banana and potato
- Upload the Trait Dictionaries in the Breeding Management System 4.0
- Release curation guidelines
- Add crops: cacao, faba bean, sunflower, forage, grape, beetroot, woody plants
- Translate the Trait Dictionaries into relevant languages

Data annotation with variables

In the Breeding Management System:

CO_322:0000684: "Estimation of Chilo partellus damage on a 1 to 5 scale"

CO_322:0000869: "Measurement of plant height in cm"

ChiloDmg_est_1to5

PH_mes_cm

In NextGen databases:

CO:0000000 CGIAR cassava trait ontology

CO:0000001 Agronomic trait

CO:0000281 Anthocyanin Pigmentation

CO:0000103 anthocyanin pigmentation visual rating 0-3

CO:0000361 Ease of Harvest

CO:0000225 ease of harvest assessment 1-3